(3) (3) (3)



AZRA NAHEED/MEDICAL COLLEGE

MBBS 4th Year (2015-16)
Pathology Revision Test (Objective)

Resource Person : Prof. Naila
Roll No: 12.205

Total Marks: 30 Time Allowed: 30 mins

Name of Student:

Sherpena Awan

Instructions:

- All objective questions are to be attempted on the paper and returned to the invigilator within specified time
 after you have received the question paper.
- Any cuttings or overwriting in answering the objective part will not be accepted and no marks will be given
 even if the answer is correct.
- 1. A 45 year old man presented with malaise, anorexia and vorniting to the emergency room. Clinically the patient had mild jaundice and liver function tests revealed normal alkaline phosphatase and raised ALT, AST. Histopathology report showed Mallory bodies in the hepatocytes. Liver damage from which of the following diseases most likely account for these findings?
 - a. Billary cirrhosis
 - b. Viral hepatitis
 - c. Alcoholic hepatitis
 - d. Hemochromatosis
 - e. Wilson's disease
- 2. 'A 65 year old male presented with complaint of right upper quadrant pain. He was diagnosed with hepatitis <u>C virus</u>, 20 years back. He also gave history of ill healthy, fatigue, fever and loss of appetite. Biopsy specimen showed presence of bile and atypical lymphocytes. Apart from CT scan and other tests, which tumor marker will support your diagnosis?
 - a. Beta HCG
 - b. Alpha feto proteins
 - c. Carcino-embryonic antigen
 - d. CA-125
 - e. CA-15
- 3. A 43 years old multigravida presented with nausea, vomiting, fever and right upper quadrant pain. (Murphy's sign is positive) Laboratory tests shows neutophilia with a left shift. What is the most likely diagnosis?
 - a. Acute cholecystitis
 - b. Carcinoma of ampulla of vater
 - c. Cholangiocarcinoma
 - d. Cholesteriolosis
 - e. Sclerozing cholangitis

Distended mucus filled gall bladder secondary to chronic cystic duct obstruction is known as: a. Hydrocele b. Mucocele c. Metaplasia d. Dysplasia · e. Cholecystitis 5. A 65 years old man developed arthritis for which he took NSAIDS for one year. After prolonged use of NSAIDS he went to local hospital due to severe epigastric pain, nausea and hemetemesis. He was advised gastric blopsy. Which of the following lesions can be presented in the biopsy specimen? a. Adenocarcinoma b. Epithelial dysplasia c. H.pylori infection d. Hyperplastic polyp -e. Acute gastritis 6. An old lady of 65 years of age comes to a hospital emergency with fever and massive hematemsis of bright red color. On physical examination she was pale, hypotensive and techypenic. Her serological tests showed HBsAg positive. The hematemesis is complication of which of the following? Hepatitis B a. Squamous cell carcinoma b. Barett's esophagus c. Esophageal varices d. Reflux oesophagitis e. Candida albicans Infection 7. A man comes to a doctor with complains of diarrhea and abdominal pain since 6 months. On physical examination abdominal tenderness and parianal fistulas were present. Occult blood test for stool was positive. Colonoscopy showed patchy mucosal edema andulceration. Microscopic examination of biopsy showed crypt abcesses, non caseating granulomas and patches of acute and chronic inflammation. Which of the following disease has these findings? a. Sarcoidosis b. Shigellosis c. Ulcerative colitis d. Crohn's disease e. Amebiasis 8. Kayser Fleisher rings in the cornea is associated with which of the following disease? a. Hepatocellular carcinoma b. Hepatic amebiasis c. Wilson's disease d. Hemochromatosis

Primary biliary cirrhosis

- 9. A lady brought her three days old infant to a hospital. The infant vomits all oral feeds. On physical examination abdomen is distended, tender and bowel sounds were reduced. Abdominal ultrasound showed marked colonic dilatation with narrowing of the sigmoid region. Biopsy of the narrowed part showed absent ganglion cells in the sub mucosa of the wall. Which of the following disease the Infant is most likely suffering from? a. Trisomy 21 b. Volvulus · c. Hirsch sprung disease d. Intussusception e. Colonic atresia 10. A student presents to an emergency with history of difficulty in swallowing both the solid and liquid food with aspiration pneumonia due to aspiration of food. On auscultation of the lungs there were crackles on the base of the right lung. A barium swallow showed marked dilatation of the esophagus above the esophageal sphincter. Biopsy specimen of the lower esophagus will most likely show the following features: a. Multinucleated cells with intranuclear inclusions
 - b. Thrombosed vascular channels
 - c. Hyperplasia
 - d. Absence of the myenteric ganglia,
 - e. Neutrophils infiltration
- 11. Paroxysmal hypertension is most typically associated with:
 - a. Pituitary adenoma
 - b. Adrenal adenoma
 - c. Thyroid adenoma
 - d. Parathyroid adenoma
 - e. Pheochromocytoma
- 12. Diabetes insipidus is associated with a lack of:
 - a. Glucocorticoids
 - b. Insulin
 - c. Thyroid hormone
 - d. Antidiuretic hormone
 - e. Growth hormone
- 13. Truncal obesity, easy bruising and osteoporosis are associated with:
 - a. Carniopharyngioma
 - b. Adrenal adenoma
 - c. Thyroid adenoma
 - d. Parathyroid adenoma
 - e. Pheochromocytoma
- 14. Features of both Hashimoto's and granulomatous thyroiditis include:

- a. Female predominance
- b. Persistent elevation of antithyroid antibodies
- c. Lymphocytic infiltration of thyroid
- d. Hyperthyroidism
- e. Hypothyroidism

)
15. Which of the following thyroid carcinomas has the best prognosis?	
a. Undifferentiated	<i>!</i>
b. Follicular	
c. Small cell	
d. Giant cell	
e. Papillary	
16. A surgeon explores a thyrold because of a "cold" nodule of the le	eft upper pole of the
thyroid. The nodule is firm, non-encapsulated, and granular. The	
hard lymph node in the adjacent internal jagular chain. The most i	ikely diagnosis is:
a. Anaplastic carcinoma	
b. Follicular adenoma	`
c. Follicular carcinoma	
d. Lymphoma	
17. Major site of Peptic Ulcer Disease is : a. Ileum	
b. Duodenum	
c. Esophagus	
d. Pancreas	
e. Cecum	
18. The granulomas of tuberculosis are differentiated from granu	ulomas of Crohn's
disease by:	, •
a. Presence of necrosis	
- b. Presence of caseation	
c. Presence of glant cells	
d. Presence of lymphocytes	
e. Presence of epithelioid cells	
19. Familial Adenomatous Polyposis is associated with mutation	n of which one of the
following target gene?	
a. MSH2	
b. MLH1	
C. APC	
d. KRAS	
e. BRAF	
20. Ulcerative colitis involves	
a. Colon and rectum only	The second secon
b. Appendix	
c. Stomach	
d. All the gastrointestinal tract	
e. Cecum only	
21. An immune mediated diarrheal disorder triggered by inges	tion of gluten
containing foods is	
a. celiac disease	
b. giardlasis	No. of the second secon
c. crohn's disease	
d. ulcerative colitis	
e. lymphoma	

.

- 22. A 45y/F had a head trauma after which she develops the complaints of excretion of large volumes of dilute urine with an inappropriately low specific gravity. Serum sodium and osmolality are increased as a result of excessive renal loss of free water, resulting in thirst and polydipsia. Which of the following disease explains this condition? a. Diabetes insispidus b. Diabetes mellitus c. Pituitary adenoma d. Prolatinoma e. Waterhouse friderichsen syndrome 23. A disease of adrenal gland causing excess production of harmone named alsosterone. Another name for this is primary hyperalsosteronism also known as a. Cushing syndrome b. Cushing disease · c. Addisons disease d. Conn's syndrome e. Sheehan syndrome 24. Which of the following hormone is secreted from posterior lobe of pituitary? a. Thyroid stimiulating hormone b. Oxytocin c. Adrenocorticotrophic hormone d. Follicular stimulating hormone
 - e. Lutenising hormone
 - 25. A girl presents with delayed puberty, absent secondary sexual characteristics, and primary amenorrhea. She also appears to be hypertensive and hypokalemia. Which of the following enzymes is increased if there is virilization of a person?
 - 11beta Hydroxylase
 - b. 17alpha hydroxylase
 - c. 21 beta hydroxylase
 - d. Pyruvate kinase
 - e. Alcohol dehydrogenase
 - 26. Which of the following thyroid cancer is associated with MEN II syndrome?
 - -a. Papillary thyroid carcinoma
 - b. Medullary thyroid cancer
 - c. Follicular thyroid carcinoma
 - d. Anaplstic thyroid carcinoma
 - e. Thyroid lymphoma
 - 27. In which of the following pairs, the hormone of endocrine glands and its primary action is mismatched.
 - a. Calcitonin-lower blood calcium level
 - b. Parathyroid-raises blood calcium level
 - c. Somatostatin-inhibits release of glucagon
 - d. Insulin-increases blood glucose level
 - e. Glucagon-increses blood glucose level

- 28. Which of the following statement is true about Pheochromocytoma?

 a. 50% extra adrenal

 b. 45% bilateral
 - · c. 10% are biologically malignant.
 - d. 10% associated with hypertension
 - e. Do not show zell ballen arrangment
 - 29. The prolactin inhibitory factor secreted by hypothalamus is
 - a. Thyrotropin releasing hormone
 - b. 5-Hydroxytryptamine
 - c. Dopamine
 - d. Acetylcholine
 - e. Progesterone
 - 30. In which of the following thyroid carcinoma, metastatic cervical lymphadenopath may be 1st sign of disease:
 - a. Papillary
 - b. Follicular
 - /c. Anaplastic
 - d. Medullary
 - e. Lymphoma